

# *The Nonperturbative S-matrix Bootstrap*

EPFL

João Penedones

Based on work with: Correia, Elias-Miró, Guerrieri, Haring, Hebbar, Homrich, Karateev, Kuhn, Marucha, Meineri, Murali Paulos, Sahoo, Toledo, van Rees, Vieira, Vuignier

Quantum Spacetime and the Renormalization Group 2023  
5/10/2023

# Outline

- Introduction

- Applications

- $2 \rightarrow 2$  Super graviton Scattering in  $d = 9, 10, 11$

- [ '21 Guerrieri, JP, Vieira ]
    - [ '22 Guerrieri, Murali, JP, Vieira ]

- a-anomaly bootstrap in 4d

- [ '22 Karateev, Marucha, JP, Sahoo ]

- Discussion

# Introduction

# The S-matrix program




Heisenberg

The "observable quantities" in  
the theory of elementary particles

1943

"Nuclear Democracy"  
'60




Chew



# Modern QFT Bootstrap

**Goal:** map out the space of QFTs and develop nonperturbative methods to compute their observables.



**Bootstrap approach:** bound the space of theories by imposing consistency conditions on physical observables.

**Strategy:** extend recent success in CFT to QFT.



# Conformal Bootstrap

correlation functions

$$\sum \text{[diagram 1]} = \sum \text{[diagram 2]}$$

# S-matrix Bootstrap

Scattering amplitudes



Flat space limit of QFT in AdS

Applications


# Some Applications

	2D phonons	with	Elias-Miró, Guerrieri, Hebbar, Vieira
	pions	with	Guerrieri, Vieira
TODAY →	super-gravitons	with	Guerrieri, Murali, Vieira
	photons	with	Meineri, Haring, Hebbar, Karateev
	massive scalars	with	Homrich, Paulos, Toledo, van Rees, Vieira
	massive fermions	with	Hebbar, Karateev
TODAY →	4d a-anomaly	with	Karateev, Marucha, Sahoo
	2d central charge	with	Correia, Karateev, Kuhn, Vaignier
	...		

- Goals:
- bound the leading Wilson coefficients of The EFT.
  - bound the interaction strength (cubic & quartic couplings)
  - bound a-anomaly of the UV CFT
  - ...

Maximal Supergravity  $d = 9, 10, 11$

# 2 → 2 Super graviton Scattering




$$\mathbb{A}_{2 \rightarrow 2} = \mathbf{R}^4 A(s, t, u).$$

↑  
pre-factor  
fixed by SUSY

crossing symmetric function of

$$\begin{cases} s = (p_1 + p_2)^2 \\ t = (p_1 - p_3)^2 \\ u = (p_1 - p_4)^2 \end{cases}$$
$$s + t + u = 0$$

# 2 → 2 Super graviton Scattering




$$\mathbb{A}_{2 \rightarrow 2} = \mathbf{R}^4 A(s, t, u).$$

↑  
pre-factor  
fixed by SUSY

crossing symmetric function of

$$\begin{cases} s = (p_1 + p_2)^2 \\ t = (p_1 - p_3)^2 \\ u = (p_1 - p_4)^2 \end{cases}$$

$$s + t + u = 0$$




Focus on charged scalar :  
[axi-dilaton in IIB]

$$T(s, t, u) \equiv s^4 A(s, t, u) = -8\pi G_N \left( \frac{s^2}{t} + \frac{s^2}{u} \right) + \dots$$

$$\frac{T(s, t, u)}{8\pi G_N} = s^4 \left( \frac{1}{stu} + \alpha \ell_P^6 + \dots \right) \equiv s^4 A(s, t, u)$$

$$16\pi G_N = (2\pi)^{d-3} \ell_P^{d-2}$$

# 2 → 2 Super graviton Scattering




$$\mathbb{A}_{2 \rightarrow 2} = \mathbf{R}^4 A(s, t, u).$$

↑  
pre-factor fixed by SUSY

crossing symmetric function of

$$\begin{cases} s = (P_1 + P_2)^2 \\ t = (P_1 - P_3)^2 \\ u = (P_1 - P_4)^2 \end{cases}$$

$s + t + u = 0$



Focus on charged scalar :  
[axi-dilaton in IIB]

$$T(s, t, u) \equiv s^4 A(s, t, u) = -8\pi G_N \left( \frac{s^2}{t} + \frac{s^2}{u} \right) + \dots$$

$$\frac{T(s, t, u)}{8\pi G_N} = s^4 \left( \frac{1}{stu} + \alpha \ell_P^6 + \dots \right) \equiv s^4 A(s, t, u)$$

$$16\pi G_N = (2\pi)^{d-3} \ell_P^{d-2}$$

What are the allowed values of  $\alpha$  ?

EFT: 
$$S = \frac{1}{(2\pi)^7 \ell_P^8} \int d^x \sqrt{-g} \left[ \mathcal{R} + \# \alpha \ell_P^6 \mathcal{R}^4 + \dots \right]$$




# Superstrings $d=10$

## IIB

$$\alpha^{\text{IIB}} = \frac{1}{2^6} E_{\frac{3}{2}}(\tau, \bar{\tau}) \geq \frac{1}{2^6} E_{\frac{3}{2}}(e^{i\pi/3}, e^{-i\pi/3}) \approx 0.1389$$


$$\tau = \chi_s + i/g_s$$



[ '97 Green, Gutperle ]

## IIA

$$\alpha^{\text{IIA}} = \frac{\zeta(3)}{32g_s^{3/2}} + g_s^{1/2} \frac{\pi^2}{96} \geq \frac{\pi^{3/2}(\zeta(3))^{1/4}}{24\sqrt{3}} \approx 0.1403$$



$$\alpha^{\text{IIB}} = \alpha^{\text{IIA}} + O\left(e^{-\frac{2\pi}{g_s}}\right).$$

M-theory  $d = 11$

$$\alpha = \frac{(2\pi)^2}{3 \cdot 2^7} \approx 0.1028$$

String theory  $d = 9$

$$\alpha = \frac{1}{2^6} \left[ \underbrace{V^{-3/7}}_{\text{related to compactification radius}} E_{3/2}(\tau, \bar{\tau}) + \frac{2\pi^2}{3} V^{4/7} \right] \geq 0.2417$$

See [ '23 Bossard, Loty ] for lower bounds in  $d = 6, 7, 8$ .

# Numerical S-matrix Bootstrap

# PRIMAL ALGORITHM

[ '17 Paulos, JP, Toledo, Van Roes, Vieira ]

1. Write amplitude ansatz obeying:  $\left\{ \begin{array}{l} \text{Lorentz invariance (+ SUSY)} \\ \text{Crossing symmetry} \\ \text{Analyticity} \\ \text{Low energy from EFT} \end{array} \right.$

2. Impose unitarity of each partial wave

3. Minimize a linear observable of the amplitude

# PRIMAL ALGORITHM

[ '17 Paulos, JP, Toledo, Van Raes, Vieira ]

1. Write amplitude ansatz obeying:

↓

# of parameters  $\sim N^2$

{

- Lorentz invariance (+ SUSY)
- Crossing symmetry
- Analyticity
- Low energy from EFT

2. Impose unitarity of each partial wave  $\rightarrow$  spin  $l \leq L$

3. Minimize a linear observable of the amplitude



4. Extrapolate  $L \rightarrow \infty$  and  $N \rightarrow \infty$

# 1. Analytic and crossing symmetric ansatz:


$$T = 8\pi G_N s^4 \left[ \underbrace{\frac{1}{stu}}_{\text{SUGRA}} + \underbrace{\sum'_{a+b+c \leq N} \beta_{(abc)} \rho^{(s)^a} \rho^{(t)^b} \rho^{(u)^c}}_{\text{"UV completion"}} \right]$$

free parameters

$s+t+u=0$   
[ $l_p=1$ ]




$$\rho(s) = \frac{1 - \sqrt{-s}}{1 + \sqrt{-s}}$$



## 2. Unitarity constraints

[ '98 Bern et al. ]



$\Rightarrow$

$$\text{Disc} A \geq s^4 \int d\text{LIPS } A \times A$$

$$\sum_{\text{two pt}} \mathbf{R}^4_{12 \rightarrow \text{two pt}} \mathbf{R}^4_{\text{two pt} \rightarrow 34} = \mathbf{R}^4_{12 \rightarrow 34} \times s^4$$

Unitarity of


$$A_{2 \rightarrow 2} = \mathbf{R}^4 A(s, t, u).$$

$\Leftrightarrow$  Unitarity of  $T = s^4 A(s, t, u)$

$$\text{Disc } T \geq \int d\text{LIPS } T \times T$$

## 2. Unitarity constraints

[98 Bern et al.]



$$\text{Disc} A \geq s^4 \int d\text{LIPS } A \times A$$

$$\sum_{\text{two pt}} \mathbb{R}^4_{12 \rightarrow \text{two pt}} \mathbb{R}^4_{\text{two pt} \rightarrow 34} = \mathbb{R}^4_{12 \rightarrow 34} \times s^4$$

Unitarity of

$$\mathbb{A}_{2 \rightarrow 2} = \mathbb{R}^4 A(s, t, u)$$

$\Leftrightarrow$  Unitarity of  $T = s^4 A(s, t, u)$

$\Rightarrow$

$$|S_\ell(s)|^2 \leq 1$$

for  $\begin{cases} \ell = 0, 2, 4, \dots, L \\ s > 0 \text{ [grid]} \end{cases}$

$$S_\ell(s) = 1 + i N_d s^{\frac{d}{2}-2} \int_{-1}^1 dz (1-z^2)^{\frac{d-4}{2}} C_\ell^{\frac{d-3}{2}}(z) T(s, z)$$

$\begin{cases} t = -\frac{s}{2}(1-z) \\ u = -\frac{s}{2}(1+z) \end{cases}$   
 $\cos \theta \approx$  scattering angle


$\swarrow$  Gegenbauer polynomial  
 $\nearrow$



3. Minimize  $\alpha$  for each  $N$  and  $L$ .


→ Semi-definite Programming SDPB

[15 Simmons-Duffin]



$d = 10$



# Positivity in the Sky



$$\left( \text{Im } T \right)_{ij} \equiv \text{Im } T \left( s, t = -\frac{s}{2}(1 - \cos \theta_{ij}), u = -\frac{s}{2}(1 + \cos \theta_{ij}) \right) \geq 0$$


This is not an independent condition but it accelerates convergence in spin ( $L \rightarrow \infty$ ).

# 4. Extrapolate $L \rightarrow \infty$ and $N \rightarrow \infty$



## Minimal values of $\alpha$


Dimension	Bootstrap	String/M-Theory
9	$0.223 \pm 0.002$	0.241752
10	$0.124 \pm 0.003$	0.138949
11	$0.101 \pm 0.005$	0.102808



**Figure 6.** The first two Regge trajectories in 10d for  $N=30$  and  $L=200$ . The error bars represent the widths and the resonances lie on curved trajectories that scale approximately like  $\ell^{1.3}$ . More details in appendix F.

Bootstrapping the  $\alpha$ -anomaly in 4d


$$d=2$$



$$c_{UV} > c_{IR}$$

[ '86 Zamolodchikov ]


$$d=4$$







$$a_{UV} > a_{IR}$$

[ '88 Cardy ]  
[ '11 Komargodski, Schwimmer ]

# S-matrix Bootstrap setup




## Mass Spectrum



$$T_{\varphi\varphi \rightarrow \varphi\varphi} = \frac{1}{f^4} a^{UV} (s^2 + t^2 + u^2) + O(s^3)$$

[11 Komargodski, Schwimmer]


# Results



**Figure 5.** Minimum possible value of the a-anomaly without any further assumptions as a function of  $1/N_{\text{max}}$  with  $L_{\text{max}} = N_{\text{max}} + 10$ . The numerical results are depicted by blue points. Linear extrapolation to  $N_{\text{max}} \rightarrow \infty$  depicted by the red line gives  $0.316 \pm 0.015$  for the minimum of  $a/a_{\text{free}}$ .

$$\frac{a}{a_{\text{FB}}} \gtrsim 0.32$$





$$N_{\max} = 20$$

$$L_{\max} = 30$$


quartic coupling

$$\lambda_0 \equiv \frac{1}{32\pi} T_{AA \rightarrow AA} (s=t=u = \frac{4}{3} m^2)$$

See [23 Marucha] for bounds without  $\mathbb{Z}_2$  symmetry.

# Discussion

- Dual formulation would be very useful.



[77' Lopez, Mennessier]

[...]

[19 Cordova, He, Kruczenski, Vieira]

[20 Guerrieri, Homrich, Vieira]

[21 Guerrieri, Elias Miro'

[21 He, Kruczenski]

[21 Guerrieri, Sever]

}  $D=2$

}  $D>2$

Unitarity:

$$2 \operatorname{Im} f_{\ell}(s) \geq |f_{\ell}(s)|^2 \Leftrightarrow \left| \frac{S_{\ell}(s)}{1 + i f_{\ell}(s)} \right|^2 \leq 1$$

vs.

Positivity:

$$\operatorname{Im} f_{\ell}(s) \geq 0$$

→ Unitarity is stronger than positivity.

Example: supergraviton scattering ( $d=10$ )

$$\begin{cases} \text{Unitarity} \Rightarrow \alpha \geq 0.14 \\ \text{positivity} \Rightarrow \alpha \geq 0 \end{cases}$$

## Future work in graviton scattering

- Other spacetime dimensions

$5 \leq d \leq 8$       Need to take into account log's from loops

$d = 4$       IR divergences

- Inelastic effects from  $2 \rightarrow 3$  and/or Black Hole production

- Other Wilson coefficients

- No SUSY.

[ '21    Bern, Kosmopoulos, Zhiboedov ]  
[ '22    Caron-Huot, Li, Parra-Martinez, Simmons-Duffin ]

# Big open questions

- Prove (or drop) Maximal Analyticity

┌ in perturbation theory? [21 Correia, Sever, Zhiboedov]  
└ from the flat space limit of QFT in AdS? [23 van Rees, Zhao]

- Anomalous thresholds
- Multi-particle amplitudes ( $n \rightarrow m$ )

Thank You!